



PCT/IL2004/000406

מדינת ישראל
STATE OF ISRAEL

REC'D 15 JUN 2004

WIPO

PCT

Ministry of Justice
Patent Office

משרד המשפטים
לשכת הפטנטים

This is to certify that
annexed hereto is a true
copy of the documents as
originally deposited with
the patent application
particulars of which are
specified on the first page
of the annex.

זאת לתעודה כי
רצופים בזה העתקים
נכונים של המסמכים
שהופקדו לכתחילה
עם הבקשה לפטנט
לפי הפרטים הרשומים
בעמוד הראשון של
הנספח.

PRIORITY DOCUMENT
SUBMITTED OR TRANSMITTED IN
COMPLIANCE WITH
RULE 17.1(a) OR (b)

This 18 MAY 2004 היום

רשם
הפטנטים
של המדינה

Commissioner of Patents



BEST AVAILABLE COPY

נתאשר
Certified

לשימוש הלישכה

For Office Use

155876

מספר:
Number

12-05-2003

תאריך:
Date

25-10-1996

הוקדם/נדחה
Ante/Post-Dated

חוק הפטנטים, תשכ"ז-1967

Patent Law, 5727 - 1967

בקשה בינלאומית לפטנט - שלב לאומי

INTERNATIONAL PATENT APPLICATION - NATIONAL PHASE

אני, (שם המבקש, מענו ולגבי גוף מאוגד - מקום התאגדותו)

I, (Name and address of applicant, and in case of body corporate-place of incorporation)

SOCIETE D'ETUDE ET DE RECHERCHE EN
INGENIERIE PHARMACUETIQUE SERIPHARM
28, Rue Sainte-Croix
F-72000 Le Mans
France

Owner, by virtue of THE LAW

בעל ההמצאה מכח הדין

of an invention the title of which is:

ששמה הוא:

תרכובות ביניים להמיסנתזה של טקסאנים ותהליכים להכנתן

(בעברית)


(Hebrew)

INTERMEDIARY COMPOUNDS FOR THE HEMISYNTHESIS OF TAXANES AND
PREPARATION PROCESSES THEREFOR

(באנגלית)

(English)

הבקשה הזאת כי ינתן לי עליה פטנט, hereby apply for a patent to be granted to me in respect thereof.

*בקשת חלוקה - Application of Division		*בקשת פטנט מוסף - Application for Patent of Addition		*דרישה דין קדימה Priority Claim	
*מבקשת פטנט from Application		*לבקשה/לפטנט to Patent/Apl.		מספר/סימן Number/Mark	תאריך Date
No. 124245 מס' מיום 25.10.1996		No. מס' מיום		95/12739	October 27, 1995
*יפוי כח: כללי / מיוחד - רצוף - בזה / עוד יוגש P.O.A.: general / individual - attached / to be filed later הוגש בענין 125049 filed in case					
		המען למסירת מסמכים בישראל Address for Service in Israel לוצאטו את לוצאטו ת.ד. 5352 באר שבע 84152			
חתימת המבקש Signature of Applicant Luzzatto & Luzzatto By:  Attorneys for Applicant DATE: 12 במאי, 2003 תאריך:		INT. APP. PCT/FR96/01676 INT. FIL. DATE: 25 October 1996 INT. PUBL. WO 97/15562		בקשה בינ"ל תאריך הגשה בינ"ל פרסום בינ"ל	

REFERENCE: 16296/03 סימוכין:

טופס זה כשהוא מוטבע בחותם לישכת הפטנטים ומושלם במספר ובתאריך ההגשה, הינו אישור להגשת הבקשה שפרטיה רשומים לעיל.

This form, impressed with the Seal of the Patent Office and indicating the number and date of filing, certifies the filing of the application the particulars of which are set out above.

מחק את המיותר Delete whatever is inapplicable

#199844

16296/03


תרכובות ביניים להמיסנתזה של טקסאנים ותהליכים להכנתן


INTERMEDIARY COMPOUNDS FOR THE HEMISYNTHESIS OF TAXANES AND
PREPARATION PROCESSES THEREFOR

This application is a divisional application from the co-pending IL 124245

The invention of IL 124245 relates to novel intermediates for the hemisynthesis of taxanes and to their processes of preparation.

Taxanes, natural substances with a diterpene skeleton which is generally esterified by a β -amino acid side chain derived from N-alkyl- or N-aroylphenylisoserine, are known as anticancer agents. Several dozen taxanes have been isolated from Taxaceae of the genus *Taxus*, such as, for example, paclitaxel ($R_1 = \text{Ac}$, $R_2 = \text{Ph}$, $R_3 = R_4 = \text{H}$), cephalomantine, their derivatives deacetylated in the 10 position, or baccatins (derivatives without side chain) represented by the formulae 1 and 2 below.





Because of concern not to rapidly exhaust its original source, *Taxus brevifolia*, French researchers have sought to isolate paclitaxel from renewable parts (leaves) of *T. baccata*, the European yew. They have thus demonstrated the probable biogenetic precursor of taxanes, 10-deacetylbaccatin III, the springboard of choice for the hemisynthesis because of its relative abundance in leaf extracts.

The hemisynthesis of taxanes, such as paclitaxel or docetaxel ($R_1 = \text{Ac}$, $R_2 = t\text{-butyloxy}$, $R_3 = R_4 = \text{H}$), thus consists in esterifying the 13-hydroxyl of a protected derivative of baccatin or of 10-deacetylbaccatin III with a β -amino acid derivative.

Various processes for the hemisynthesis of paclitaxel or of docetaxel are described in the state of the art (EP-0 253 738, EP-0 336 840, EP-0 336 841 and IL 89831, EP-0 495 713, WO 92/09529, WO 94/07877, WO 94/07878, WO 94/07879, WO 94/10169, WO 94/12482, EP-0 400 971 and IL 94426,

EP-0 428 376, WO 94/14787). Two recent works [I. Georg, T.T. Chen, I Ojima, and D.M. Vyas, "Taxane Anticancer Agents, Basic Science and Current Status", ACS Symposium Series 583, Washington (1995)] and especially
5 [Matthew Suffness, "Taxol[®] Science and Applications" CRC Press (1995) and 1500 references cited] comprise exhaustive compilations of hemisyntheses of taxanes.

The β -amino acid side chains derived from N-alkyl- or N-aroylphenylisoserine of paclitaxel or
10 docetaxel are of (2R,3S) configuration and one of the main difficulties in the hemisynthesis of taxanes is to obtain an enantiomerically pure product. The first problem consists in obtaining a pure enantiomer of the phenylisoserine derivatives employed in the
15 hemisynthesis of taxanes. The second problem consists in retaining this enantiomeric purity during the esterification of the baccatin derivative and the subsequent treatments of the products obtained (deprotection of the hydroxyls, and the like).

20 Many studies on asymmetric synthesis involving derivatives of β -amino acids have focused on the chemistry of isoserine and of its derivatives, β -amino acids for which a dehydrated cyclic form is a β -lactam (EP-0 525 589). The majority of the various
25 syntheses of phenylisoserine derivatives useful as precursors of taxane side chains focus on a common intermediate, (2R,3R)-cis- β -phenylglycidic acid, which is subsequently converted to β -phenylisoserine by

reaction with ammonia (EP-0 495 718) or a nucleophile (Gou et al., *J. Org. Chem.*, 1983, 58, 1287-89). These various processes require a large number of stages in order to produce β -phenylisoserine of (2R,3S)

5 configuration, necessarily with a stage of racemic resolution by conventional selective crystallization techniques, either for cis- β -phenylglycidic acid or for β -phenylisoserine, or subsequently, after conversion.

Furthermore, in order to retain the enantiomeric purity

10 of taxane side chain precursors during the esterification of the baccatin derivative, various means have been provided, in particular by using cyclic intermediates of blocked configuration, which remove the risks of isomerization during esterification

15 reactions under severe reaction conditions. In particular, they involve β -lactam (EP-0400 971 and IL89831), oxazolidine (WO 92/09539, WO 94/07877, WO 94/07878, WO 94/07879, WO 94/10169, WO 94/12482), oxazinone (EP-0 428 376) or oxazoline (WO 94/14787) derivatives.

20 These cyclic precursors are prepared from the corresponding β -phenylisoserine derivative. As for the latter, the processes provided involve a large number of stages and a necessary racemic resolution in order to obtain the desired taxane side chain precursor. It


25 was thus important to develop a novel route for the improved synthesis of intermediates which are taxane side chain precursors, in particular of enantiomers of cis- β -phenylglycidic acid, of β -phenylisoserine and of

their cyclic derivatives.

Finally, for the hemisynthesis of taxanes and in particular of paclitaxel, the sole appropriate baccatin derivative used until now is that for which the 7-hydroxy radical is protected by a trialkylsilyl (EP-0 336 840, WO 94/14787), the deprotection of which is carried out exclusively in acidic medium. It was thus also important to employ novel protective groups for the hydroxyl functional group which in particular make possible selective protection of the 7-hydroxy radical and in addition allow a wider choice of operating conditions for the deprotection stage.

The invention of IL 124245 relates first of all to an improved process for the preparation of taxane side chain precursors.

The process according to the invention of IL 124245 consists in converting a cis- β -arylglycidate derivative of general formula I




in which

Ar represents an aryl, in particular phenyl, and R represents a hydrocarbon radical, preferably a linear or branched alkyl or a cycloalkyl optionally substituted by one or more alkyl groups,

so as to regio- and stereospecifically introduce the

β -N-alkylamide and the α -hydroxyl or their cyclic precursors in a single stage by a Ritter reaction. Depending on the reaction mixture, two types of Ritter reaction are thus distinguished: one with opening of the oxetane, resulting in a linear form of the chain which is directly and completely functionalized, the other resulting in the direct formation of an oxazoline. The "*" symbol indicates the presence of an asymmetric carbon, with an R or S configuration. In both cases, the Ritter reaction is stereospecific, with retention of C-2 configuration and inversion of C-3 configuration. The process according to the invention is advantageously carried out on one of the enantiomers of the cis- β -arylglycidate derivative of general formula I, so as to obtain the corresponding enantiomer of the linear chain or of the oxazoline obtained, without subsequently requiring a racemic resolution. According to the method of preparation of the cis- β -arylglycidate derivative of general formula I described subsequently, R represents an optically pure enantiomer of a highly sterically hindered chiral hydrocarbon radical, advantageously a cycloalkyl substituted by one or more alkyl groups, in particular a cyclohexyl. R will then preferably be one of the enantiomers of the menthyl radical, in particular (+) menthyl.

The present invention relates to a process for the preparation of taxane side chain precursors in which a cis- β -arylglycidate derivative of general formula I



5 in which

Ar represents an aryl radical and
R represents an optically pure enantiomer of
a highly sterically hindered chiral hydrocarbon
radical, preferably a branched alkyl or a cycloalkyl
10 optionally substituted by one or more alkyl groups,
is converted, so as to regio- and stereospecifically
introduce the β -N-alkylamide and the α -hydroxyl or their
cyclic precursors in a single stage by a Ritter
reaction, which consists:

15 of the direct synthesis of a cyclic chain by
reacting a cis- β -arylglycidate derivative of
general formula I defined above with a nitrile of
formula




20

in which

R', represents an aryl radical or a lower alkyl or
lower perhaloalkyl radical, such as
trichloromethyl,


in the presence of a Lewis acid or of a protonic acid, in anhydrous medium, in order to obtain the oxazoline of general formula IIb



in which Ar, R and R', are defined above.

5

A preferred embodiment of the process of this invention is characterized in that the cis-β-arylglycidate derivative of general formula I




10

in which

Ar is defined above and


R represents an optically pure enantiomer of a highly sterically hindered chiral hydrocarbon radical, preferably a branched alkyl or a cycloalkyl optionally substituted by one or more alkyl groups,

is prepared by reacting the aldehyde of formula



15

with the haloacetate of formula



Ar and R being defined above and

X representing a halogen, in particular a chlorine or a bromine.

In another preferred embodiment of the invention there is provided a process characterized in that R represents an optically pure enantiomer of a highly sterically hindered chiral hydrocarbon radical, advantageously a cycloalkyl substituted by one or more alkyl groups, in particular a cyclohexyl.


Accordingly, R is one of the enantiomers of the menthyl radical, in particular (+)-menthyl.

In a further preferred embodiment, the process according to the invention is characterized in that Ar and R₂ represent a phenyl.

The Lewis acid is chosen from the boron trifluoride acetic acid complex, boron trifluoride etherate, antimony pentachloride, tin tetrachloride or titanium tetrachloride and the protonic acid is tetrafluoroboric acid.

Still a further preferred embodiment of a process, the invention is characterized in that the derivatives of formula IIb defined as in claim 1 in which R represents a hydrogen atom are obtained by controlled saponification.

This invention further provides a compound of formula:



IIb

in which:

Ar represents an aryl radical,


R represents an optically pure enantiomer of
a highly sterically hindered chiral hydrocarbon
radical, preferably a branched alkyl or a cycloalkyl
optionally substituted by one or more alkyl
groups and

R', represents aryl radical above or a lower
alkyl or lower perhaloalkyl radical, such
as trichloromethyl.

1. Direct synthesis of the linear chain

The direct synthesis of the linear chain by

the Ritter reaction consists in reacting a
 cis- β -arylglycidate derivative of general formula I
 defined above with a nitrile of formula




5 in which

R_2 represents an aryl radical, preferably a
 phenyl,

in the presence of a proton acid, such as sulphuric
 acid, perchloric acid, tetrafluoroboric acid, and the
 10 like, and of water.

A β -arylisoserine derivative of general
 formula IIa



in which Ar, R and R_2 are defined above,
 is then obtained.

15 The reaction is carried out with inversion of
 the configuration of the C-3 of the
 cis- β -phenylglycidate derivative. Thus, starting from a
 (2R,3R)-cis- β -phenylglycidate derivative, the
 corresponding β -arylisoserine derivative of (2R,3S)
 20 configuration is obtained.

The Ritter reaction is carried out in an
 appropriate solvent, at a temperature of between -75
 and +25°C.


The appropriate solvent can be the nitrile

itself, when it is liquid at the reaction temperature, or alternatively the acid itself (sulphuric ac., perchloric ac. or tetrafluoroboric ac.), or a solvent, such as, for example, methylene chloride or ethyl ether. The proton acids conventionally used can contain the water necessary for the hydrolysis.

When benzonitrile (R_2 = phenyl) is employed with the cis- β -arylglycidate of general formula I of (2R,3R) configuration for which Ar represents a phenyl, then the corresponding β -arylserine derivative of general formula IIa of (2R,3S) configuration for which Ar and R_2 represent a phenyl is directly obtained, which product is none other than the precursor of the side chain of paclitaxel.

2. Direct synthesis of the cyclic chain

For this second possibility, a Ritter reaction is also carried out with a nitrile of formula



in which


R'_2 represents R_2 defined above or a lower alkyl or lower perhaloalkyl radical, such as trichloromethyl,

in the presence of a Lewis acid, in particular the boron trifluoride acetic acid complex, boron trifluoride etherate, antimony pentachloride, tin tetrachloride, titanium tetrachloride, and the like, or of a proton acid, such as, for example, tetrafluoroboric acid, the reaction being carried out

in anhydrous medium.

As for the synthesis of the linear chain, the solvent can be the nitrile itself, when it is liquid at the reaction temperature, or alternatively an appropriate solvent, such as, for example, methylene chloride or ethyl ether. The reaction temperature is also between -75 and $+25^{\circ}\text{C}$.

In the absence of water, an intramolecular Ritter reaction is carried out and the oxazoline of general formula IIb




in which Ar, R and R' , are as defined above, is obtained.

As in the Ritter reaction in the presence of water, the reaction is carried out with inversion of the configuration of the C-3 of the cis- β -phenylglycidate derivative. Thus, starting from a (2R,3R)-cis- β -phenylglycidate derivative, the corresponding oxazoline of (2R, 3S) configuration is obtained.

For both Ritter reactions, in order to avoid the formation of a free carbocation which is the cause of many potential side reactions, the reactants are preferably added in the following order: i) the complex between the nitrile and the acid is first formed, then

ii) the acid catalyst is added to the mixture composed of the oxirane and the nitrile.

The products obtained by this first stage, which are β -arylisoserine derivatives of general formula IIa or oxazoline derivatives of general formula IIb, can be further converted in a second optional stage described hereinbelow or then converted to acids by controlled saponification, before being coupled to a protected baccatin derivative for the hemisynthesis of taxanes, in particular of paclitaxel and its 10-deacetylated derivatives or of docetaxel. In the case of β -arylisoserine derivatives of general formula IIa, the saponification can be preceded by a conventional stage of protection of the hydroxyl by an appropriate protective group. A derivative of general formula II'a



in which

Ar, R and R_2 are defined above, and GP represents a protective group for the hydroxyl functional group which is appropriate for the synthesis of taxanes, in particular chosen from alkoxy ether, aralkoxy ether, aryloxy ether or haloalkoxycarbonyl radicals, such as, for example, methoxymethyl, 1-ethoxyethyl, benzyloxymethyl or

(β -trimethylsilylethoxy)methyl groups, tetrahydropyranyl or β -alkoxycarbonyl (TrOC) radicals, β -halogenated or alkylsilyl ethers or alkoxyacetyl, aryloxyacetyl, haloacetyl or formyl radicals, is then obtained.

3. Possible conversion of the derivatives of formula IIa or IIb


The derivatives of general formula IIa or IIb obtained above can optionally be converted into novel intermediates which are side chain precursors in the hemisynthesis of taxanes. These conversions take place with retention of the configuration of the C-2 and C-3 positions. The novel intermediates obtained will thus have the same stereochemistry as the derivatives of formula IIa or IIb from which they derive. The products obtained in this second stage are subsequently converted into acids by controlled saponification, before being coupled with a protected baccatin derivative for the hemisynthesis of taxane, in particular of paclitaxel or of docetaxel.

3.1 Cyclization of the derivatives of general formula IIa

The derivatives of general formula IIa can subsequently be converted into oxazolines of formula IIb according to conventional methods of the state of the art (WO 94/14787).


The β -arylisoserine derivatives of general formula IIa can also be converted into novel

oxazolidinone cyclic intermediates of general formula
III'a



in which Ar and R are defined above and R''₂ represents
R'₂, defined above, an alkoxy radical, preferably a
5 t-butoxy radical, or a linear or branched alkyl radical
comprising at least one unsaturation, for example a
1-methyl-1-propylene radical, and the corresponding
dialkyl acetals.

The oxazolidinones of general formula III'a
10 are obtained first of all by reacting a β-arylisoserine
derivative of general formula IIa with a
haloalkoxycarbonyl ester, in particular 2,2,2-
trichloroethoxycarbonyl (TroC), and then by cyclization
in the presence of a strong organic base, such as
15 diazabicycloundecene (DBU). An oxazolidinone derivative
of general formula IIIa



in which Ar and R are defined above,
is then obtained.

The derivatives of general formula IIIa can
20 also be obtained by direct synthesis, by reacting the


β -arylglycidate derivatives of formula II'a with urea.

The acylated derivatives of general formula III'a are obtained by introducing the R''_2 -CO- radical according to the usual acylation techniques, in the presence of an appropriate acylating agent, for example an acyl halide of formula R''_2 -CO-X, in which R''_2 is defined above and X represents a halogen, or an anhydride of the corresponding acid.

The dialkyl acetals are obtained according to the usual techniques for the formation of acetals.

3.2 Opening of the oxazoline of general formula IIb


The β -arylisoserine derivative of general formula IIb



in which Ar, R and R'_2 are defined above, is obtained by hydrolysis of the oxazoline of general formula IIb in acidic medium.

Advantageously, when R'_2 represents a lower perhaloalkyl, such as trichloromethyl, the R'_2 -CO- radical constitutes a protective group for the hydroxyl functional group.

This taxane side chain precursor can then be converted into amides of general formula III'b




in which .

Ar, R, R', and R'' are defined above.

The precursor of the side chain of paclitaxel (R'' = phenyl) or of docetaxel (R'' = t-butoxy) can thus be obtained without distinction.

4. Preparation of the cis-β-arylglycidic acid derivative of formula I

The cis-β-arylglycidic acid derivative of formula I can be prepared according to conventional processes of the state of the art or by simple esterification of cis-β-arylglycidic acid with the corresponding alcohol R-OH. In order to improve the overall yield in the synthesis of taxane chain precursors, a cis-β-arylglycidate derivative of general formula I



in which

Ar is defined above and

R represents an optically pure enantiomer of a highly sterically hindered chiral hydrocarbon


radical,

is prepared in the process according to the invention

by reacting the aldehyde of formula



with the haloacetate of formula



5 Ar and R being defined above and
X representing a halogen, in particular a chlorine
or a bromine.

Advantageously, the optically pure enantiomer
of a highly sterically hindered chiral hydrocarbon
10 radical is a cycloalkyl substituted by one or more
alkyl groups, in particular a cyclohexyl.

It concerns a Darzens' reaction through which
a mixture of the two diastereoisomers, ester of
(2R,3R)-cis- β -arylglycidic acid and (2S,3S)-cis- β -
15 arylglycidic acid and of an optically pure enantiomer
of the chiral alcohol R-OH, is obtained, since the
Darzens' reaction, carried out with a highly sterically
hindered haloacetate, results essentially in the cis
form of the β -arylglycidate. Advantageously, the highly
20 sterically hindered chiral hydrocarbon radical will be
chosen so that it allows the physical separation of the
two diastereoisomers from the reaction mixture, for
example by selective crystallization, without requiring
a stereospecific separation of the desired enantiomer
25 at the end of the reaction by conventional
crystallization or chiral column chromatography
methods.

Advantageously, R-OH represents menthol, one

of the rare highly sterically hindered chiral alcohols which is economic and commercially available in both its enantiomeric forms.

In the process for the synthesis of a precursor of the taxane side chain, the goal is to prepare a cis- β -phenylglycidate of (2R,3R) configuration. In this case, the highly sterically hindered chiral hydrocarbon radical R will be selected so that the diastereoisomer of the cis- β -phenylglycidate of (2R,3R) configuration crystallizes first from the reaction mixture. When R-OH is menthol, (+)-menthol is advantageously employed.

The asymmetric Darzens' reaction is carried out in the presence of a base, particularly an alkali metal alkoxide, such as potassium tert-butoxide, or an amide, such as lithium bis(trimethylsilyl)amide, in an appropriate solvent, in particular an ether, such as ethyl ether, at a temperature of between -78°C and 25°C. The reaction results in a diastereoisomeric mixture composed virtually exclusively of the cis-glycidates, which can reach a yield of greater than 95%, in the region of 97%. Treatment of the isolated product in an appropriate solvent, in particular a methanol/water mixture, makes it possible readily to obtain physical separation of the required diastereoisomers.

By fractional crystallization (2 stages), rapid enrichment in the desired diastereoisomer is

obtained, with a diastereoisomeric purity of greater than 99%.

The latter point is particularly important because it conditions the isomeric purity of the final taxane, the undesirable diastereoisomers exhibiting their own biological activity which is different from that of the desired taxane.

It is remarkable to observe that the selective use of the two enantiomers of the menthyl ester makes it possible to access, using the same process, the 2 precursor diastereoisomers of the two enantiomers of glycidic acid.

In addition to a fairly high yield of pure isolated diastereoisomer (up to 45%), the diastereoisomeric purity of the major product of the reaction, the ease of implementation of the reaction, the simplicity and the speed of the purification, and the low cost of the reactants and catalysts make the industrial synthesis of this key intermediate in the asymmetric synthesis of β -amino acids easy and economical to access.

When a derivative of general formula I obtained by an asymmetric Darzens' reaction is used in the process according to the invention, the derivatives of general formulae IIa, II'a, IIb, IIIa, IIIb and III'b defined above are then obtained for which R represents an optically pure enantiomer of a highly sterically hindered chiral hydrocarbon radical, such as

a cycloalkyl substituted by one or more alkyl groups, in particular a cyclohexyl, preferably menthyl, advantageously (+)-menthyl.

5 The present invention also relates to these derivatives, which are of use as intermediates in the synthesis of taxane side chains.

It should be noted that the present process constitutes a very rapid access to the substituted chiral oxazolines already described in the literature
10 (WO 94/14787), in 3 stages from commercially available products instead of 6 to 8.

5. Controlled saponification

A controlled saponification of the derivatives of general formulae IIa, II'a, IIb, IIIa, IIIb and III'b is carried out under mild conditions, so
15 as to release the acidic functional group while retaining the structure of the said derivatives, for example in the presence of an alkali metal carbonate in a methanol/water mixture.

20 After controlled saponification, the derivatives of general formulae IIa, II'a, IIb, IIIa, IIIb and III'b defined above for which R represents a hydrogen atom are obtained, which derivatives can be employed directly in the hemisynthesis of taxanes by
25 coupling with an appropriate baccatin III derivative.

6. Hemisynthesis of taxanes

6.1 Esterification

The present invention thus also relates to a








process for the hemisynthesis of taxanes of general formula IV,

C-B

IV

in which

5 C represents a side chain chosen from the radicals of following formulae:



in which Ar, R_2 , R'_2 , R''_2 , R_3 and GP are defined above, and

B represents a radical derived from baccatin III of general formula V

Ac represents the acetyl radical,

Bz represents the benzoyl radical,

Me represents the methyl radical,

5 R_1 represents an acetyl radical or a protective group for the hydroxyl functional group GP1, and R_2 represents a protective group for the hydroxyl functional group GP2.

by esterification of an appropriate baccatin III derivative of general formula V, carrying a C-13 hydroxyl functional group, with one of the derivatives of general formulae IIa, II'a, IIb, IIIa, III'a, IIIB and III'b defined above, for which R represents a hydrogen atom, under conventional conditions for the preparation of taxanes as defined in the state of the art (in particular: EP-0 253 738, EP-0 336 840, EP-0 336 841 and IL 89831, EP-0 495 718, WO 92/09589, WO 94/07877, WO 94/07878, WO 94/07879, WO 94/10169, WO 94/12482, EP-0 400 971 and IL 94426, EP-0 428 376, WO 94/14787).

The GP1 and GP2 protective groups are, independently of one another, conventional groups employed in the hemisynthesis of taxanes, such as trialkylsilyls (EP-0 336 840) or TrOC (EP-0 336 841).

5 GP1 and GP2 also represent, independently of one another, linear or branched hindered haloalkoxycarbonyl radicals comprising at least one halogen atom. They are advantageously radicals in which the alkyl residue comprises between 1 and 4 carbon
10 atoms and 3 or 4 halogen atoms, preferably chosen from 2,2,2-tribromoethoxycarbonyl, 2,2,2,1-tetrachloroethoxycarbonyl, 2,2,2-trichloro-t-butoxycarbonyl and trichloromethoxycarbonyl radicals, radicals which are all more hindered than the haloalkoxycarbonyl (TrOC)
15 used until now to protect taxanes in the 7 position.


GP1 and GP2 also represent, independently of one another, acyl radicals in which the carbon α to the carbonyl functional group carries at least one oxygen atom.

20 These acyl radicals are described in particular in Patent Application EP-0 445 021. They are advantageously alkoxy- or aryloxyacetyl radicals of formula



25 in which R_6 represents a sterically hindered alkyl radical, a cycloalkyl radical or an aryl radical, or arylidenedioxyacetyl radicals of formula

in which Ar[•]
represents an
arylidene radical.



5 Sterically hindered alkyl is preferably
understood to mean a linear or branched C₁-C₆ alkyl
radical substituted by one or more bulky substituents
chosen from halogens or linear or branched C₁-C₆ alkyl,
linear or branched C₁-C₆ alkoxy or C₁-C₆ cycloalkyl or
aryl radicals. It will be, for example, a tert-butyl or
10 triphenylmethyl radical.


Cycloalkyl is preferably understood to mean a
C₃-C₆ cycloalkyl radical optionally substituted by one
or more bulky substituents chosen from halogens or
linear or branched C₁-C₆ alkyl, linear or branched C₁-C₆
15 alkoxy or aryl radicals. Advantageously, it is a
cyclohexyl radical substituted by one or more linear or
branched C₁-C₆ alkyl radicals, such as, for example,
menthyl, its racemate or its enantiomers and their
mixtures in all proportions.

20 Aryl is preferably understood to mean a
phenyl, naphthyl, anthryl or phenantryl radical
optionally substituted by one or more bulky
substituents chosen from halogens or linear or branched
C₁-C₆ alkyl, linear or branched C₁-C₆ alkoxy or aryl
25 radicals, in particular the phenyl radical. It is
preferably a phenyl radical optionally substituted by
one or two above bulky substituents ortho- and ortho'-

to the ether bond.

Finally, arylidene is preferably understood to mean a phenylene, naphthylene, anthrylene or phenanthrylene radical optionally substituted by one or more bulky substituents chosen from halogens or linear or branched C₁-C₆ alkyl, linear or branched C₁-C₆ alkoxy or aryl radicals, in particular the phenyl radical.

GP1 and GP2 also represent, independently of one another, a trialkylgermanyl radical or together form a divalent radical of formula




in which

R₇ and R₈, independently of one another, represent a sterically hindered alkyl radical as defined above; in particular, R₇ and R₈ each represent an isopropyl radical.

6.2 Optional opening

When C represents a radical of formula IIb or IIIa, the oxazoline ring is opened in order to obtain a taxane derivative of formula VI



in which

Ac, Bz, Me, Ar, R₂, R₄ and R₅ are defined above.

The IIb, IIIa and III'a radicals are generally opened by hydrolysis in acidic or basic medium. The radical of formula IIb can be opened according to the methods described in the state of the art (in particular WO 94/14787), by hydrolysis in acidic medium, followed by treatment in basic medium, in order to obtain the derivative of general formula VI.

6.3 Deprotection

Finally, the hydroxyls of the derivatives of general formula V or VI are deprotected by replacing the protective groups for the hydroxyl functional group, GP (when C represents the II'a radical), GP1 (when R₄ is other than an acetyl) and GP2, by a hydrogen atom according to the usual techniques.

For the derivatives of general formula V in which C represents a radical of formula IIb or IIIa and

GP1 and/or GP2 are, independently of one another, conventional groups employed in the hemisynthesis of taxanes, such as trialkylsilyls, the deprotection is carried out simultaneously with the opening described above.

When GP1 and/or GP2 are bulky haloalkoxycarbonyl radicals, deprotection is carried out according to the usual techniques described for TrOC, by the action of zinc or of zinc doped with heavy metals, such as copper, in an organic solvent, in particular in acetic acid, tetrahydrofuran or ethyl alcohol, with or without water.

When GP1 and/or GP2 are acyl radicals in which the carbon α to the carbonyl functional group carries at least one oxygen atom, deprotection is carried out in basic medium by saponification in methanol at low temperature, advantageously with ammonia in methanol at a temperature of less than 10°C, preferably in the region of 0°C.

For the case where C represents a radical of formula IIb, opening of the oxazoline is carried out simultaneously with deprotection in basic medium, in order to result, in one stage, in the corresponding taxane derivative of general formula VI in which R_4 represents an acetyl radical or a hydrogen atom and R_5 represents a hydrogen atom, in contrast to the opening in acidic medium described in the state of the art,

which requires a second stage in basic medium.

The known protective groups are removed using known methods and the oxazoline chain, when it was present, opened out by hydrolysis, giving taxanes in every respect identical to the reference taxanes. By way of example, and in order to show the validity of the invention without, however, limiting the scope thereof, paclitaxel, 10-deacetyltaxol, cephalomanine and docetaxel can be obtained from the corresponding protected derivatives.

The deblocking of the acyls in which the carbon α to the carbonyl functional group carries at least one oxygen atom was first attempted under the conventional conditions regarded as the mildest, that is to say zinc acetate in methanolic medium at reflux. In this case, the reaction being complete in a few hours (against a few days for acetates), we constantly isolated, in addition to the desired product, its C-7 epimer resulting from the conventional retroaldolization equilibrium. It being presumed that, even under the neutral, indeed slightly acidic, conditions, the main agents responsible were methanol and especially the temperature, we returned to the standard conditions for deblocking acyls described by early writers, by saponification in basic medium in ethanol at low temperature. Under these conditions, no significant epimerization was observed. By way of example, we obtained paclitaxel, 10-deacetyltaxol,

cephalomanine and docetaxel, in every respect identical to the reference taxanes, from the corresponding alkoxy- or aryloxyacetylated derivatives.

Finally, it should be noted that all the methods described previously, which are nevertheless targeted at improving the overall yield of the hemisynthesis, consist in synthesizing the phenylisoserine chain beforehand, for the purpose of converting it into one of the cyclic structures mentioned above (β -lactams, oxazolidines or oxazolines). Thus, paradoxically, the apparent better performances in the coupling of these cyclic structures only compensates for the fall in overall yield caused by the addition of ring creation stages to the synthetic sequence for the linear chain (i.e. a total of 9 stages). For the general process for the synthesis of taxanes according to the invention, a product such as paclitaxel is obtained in only 5 stages:

- (1S,2R,5S) - (+) -menthyl (2R,3R) -3-phenylglycidate
- (1S,2R,5S) - (+) -menthyl (4S,5R) -2,4-diphenyl-4,5-dihydroazole-5-carboxylate
- saponification
- hemisynthesis (esterification)
- opening and deprotection.

Finally, the present invention relates to the synthetic intermediates of general formulae IV, V and VI described above which are of use in the general synthesis of taxanes, a subject of the present

invention.

Generally, hydroxycarbon radical is preferably understood to mean according to the invention a saturated or unsaturated hydrocarbon radical which can comprise one or more saturations, such as an optionally unsaturated linear or branched alkyl, an optionally unsaturated cycloalkyl, an aralkyl or an aryl, it being possible for each optionally to be substituted by one or more substituents, in particular alkyl substituents.

Linear or branched alkyl is preferably understood to mean according to the invention a C_1 - C_6 alkyl, in particular chosen from the methyl radical, ethyl radical, propyl radical, isopropyl radical, butyl radical and its various branched isomers, such as, for example, tert-butyl, pentyl radical and hexyl radical and their various branched isomers. This definition also applies to the alkyl residues of the alkoxy or aralkoxy radicals.

Cycloalkyl is preferably understood to mean according to the invention a C_3 - C_6 cycloalkyl, in particular chosen from the cyclopropyl, cyclobutyl, cyclopentyl or cyclohexyl radicals.

Aryl is preferably understood to mean according to the invention an aromatic or heteroaromatic radical, in particular chosen from the phenyl, naphthyl, anthryl, phenantryl, pyridyl or pyrimidyl radicals and the like.


Finally, halogen is preferably understood to mean chlorine, bromine or iodine. The haloalkoxy-carbonyl radicals are preferably radicals in which the alkyl residue comprises between 1 and 4 carbon atoms and 3 or 4 halogen atoms.



The general process for the synthesis of taxanes according to the invention is repeated in Scheme 1 below, for R representing (+)-menthyl and R₂ or R'₂ representing phenyl.

The final stage in the hemisynthesis of taxanes by the process according to the invention is summarized in Schemes 2 and 3 below. Scheme 2 summarizes the synthesis of paclitaxel from derivatives of formula IV defined above in which C represents a radical of formulae IIb or III'a. Scheme 3 summarizes the synthesis of 10-deacetyltaxol from a derivative of formula IV in which C represents a radical of formula IIb.

Of course, the same synthetic schemes can be used for the other definitions of the substituents.

Material which is outside the scope of the claims does not constitute a part of the claimed invention.



SCHEME 2SCHEME 3


GP₁, GP₂ = triethylgermanium, 2,2,2-trichloro-t-butoxycarbonyl

EXPERIMENTAL PART

I. Taxane side chain precursors

Example 1:

(1S,2R,5S) - (+) -Menthyl chloroacetate




5 57 mL (0.704 mol) of anhydrous pyridine are
 added to a stirred solution at room temperature of
 100 g (0.640 mol) of (1S,2R,5S) - (+) -menthol in 1 L of
 dry dichloromethane. After stirring for a few minutes,
 56 mL (0.704 mol) of chloroacetyl chloride are
 10 subsequently added and the reaction is allowed to
 continue for 30 min. After monitoring by T.L.C., 50 g
 of crushed ice are added and the reaction mixture is
 left vigorously stirring for 1 h. After diluting with
 100 mL of dichloromethane, the organic phase is washed
 15 several times with a saturated aqueous sodium chloride
 solution (200 mL), dried over MgSO_4 and then
 concentrated under reduced pressure. After purifying
 the crude product thus obtained by silica gel
 chromatography (15-40 μm) (eluent: cyclohexane/ethyl
 20 acetate, 20/1), 146 g of (1S,2R,5S) - (+) -menthyl
 chloroacetate are obtained in the form of a syrup.

The compound obtained exhibits the following
 characteristics:

• 400 MHz ^1H NMR (CDCl_3) (δ ppm): 4.77 (1H, dt), 4.06 and 4.02 (2H, 2d, $J = 13.6$ Hz), 2.02 (1H, m, $J = 11.8$ Hz), 1.87 (1H, m, $J = 7$ and 2.6 Hz), 1.69 (2H, m), 1.50 (1H, m), 1.43 (1H, m, $J = 11.7$ and 3 Hz), 1.07 (1H, m), 1.02 (1H, q, $J = 11.8$ Hz), 0.92 and 0.90 (6H, 2d, $J = 6.4$ Hz), 0.89 (1H, m), 0.77 (3H, d, $J = 7$ Hz).

Example 2:

(1S,2R,5S)-(+)-Menthyl (2R,3R)-3-phenyl-glycidate



69 mL (0.686 mol) of benzaldehyde are added to a stirred solution at room temperature of 152 g (0.653 mol) of (1S,2R,5S)-(+)-menthyl chloroacetate in 600 mL of anhydrous ethyl ether. After stirring for a few minutes, the solution is cooled to -78°C under an inert atmosphere, a suspension of 85 g (0.718 mol) of potassium tert-butoxide in 400 mL of anhydrous ethyl ether is subsequently added over 2 h and the reaction mixture is allowed to return to room temperature. After monitoring by T.L.C., the organic part is diluted with 200 mL of dichloromethane, washed several times with a saturated sodium chloride solution, dried over MgSO_4 and concentrated under reduced pressure. 200 g of a crude product are thus obtained in the form of a syrup

containing four diastereoisomers (of which two are *cis* and two are *trans*), which is subjected as is to a fractional crystallization.

In a first step, the solution of the crude product in 2 L of methanol brought to 60°C, to which 700 mL of osmosed water are gradually added, is left for 16 h at room temperature without being subjected to vibrations. A yellow-coloured lower solid phase rich in *trans* isomers is discarded and the white crystals of the upper phase, which are rich in *cis* isomers, are separated by filtration. The crystals thus obtained are redissolved in 2 L of methanol brought to 60°C, 500 mL of osmosed water are added, until a persistent cloudiness is obtained, and the mixture is left for 16 h at room temperature. Three additional crystallizations, carried out according to the same process but with reduced volumes of methanol (1 L) and water (200 mL), are necessary to obtain 23 g of (1*S*,2*R*,5*S*)-(+) -menthyl (2*R*,3*R*)-3-phenylglycidate in the crystalline state with an HPLC purity > 99% (Yd = 12%).

The compound obtained exhibits the following characteristics:

- M.p. = 104°C
- 400 MHz ¹H NMR (CDCl₃) (δ ppm): 7.40 (2H, dd, J = 7.8 Hz and 1.7 Hz), 7.32 (3H, m), 4.58 (1H, dt, J = 10.9 Hz and 4.2 Hz), 4.26 (1H, d, J = 4.6 Hz), 3.83 (1H, d, J = 4.8 Hz), 1.6 to 0.85 (9H, m), 0.78 (3H, d, J = 7 Hz), 0.75 (3H, d, J = 6.4 Hz), 0.62 (3H, d,

$J = 6.9 \text{ Hz}$).

X-ray diffraction of a (1*S*,2*R*,5*S*)-(+) -menthyl (2*R*,3*R*)-3-phenylglycidate single crystal for the purpose of the indirect determination of the absolute configuration:

5 The single crystal was obtained from a crystalline suspension resulting from the addition, while hot, of the non-solvent (water) to a semi-saturated solution of the glycidate in methanol. On slow cooling, fine needles with a purity of 99.95%
10 (HPLC) were deposited by this solution, which needles were stored under moist conditions until the final selection.

 The selected sample (fine needle with dimensions 0.12 × 0.12 × 0.40 mm) was studied on a CAD4
15 Enraf-Nonius automatic diffractometer (molybdenum radiation with graphite monochromator). The unit-cell parameters were obtained by refinement of a set of 25 reflections with a high theta angle. Data collection ($2\theta_{\text{max}} = 50^\circ$, scanning $\omega/2\theta = 1$, $t_{\text{max}} = 60 \text{ s}$, HKL domain:
20 H 0.6 K 0.14 L 0.28, intensity controls without significant drift (0.1%)) provided 1888 reflections, 1037 of which with $I > 1.5\sigma(I)$.

$\text{C}_{19}\text{H}_{26}\text{O}_3$: Mr = 302.42, orthorhombic, $P2_12_12_1$,
 $a = 5.709(11)$, $b = 12.908(4)$, $c = 24.433(8) \text{ \AA}$,
25 $V = 1801(5) \text{ \AA}^3$, $Z = 4$, $D_s = 1.115 \text{ Mg.m}^{-3}$, $\lambda(\text{MoK}\alpha) = 0.70926 \text{ \AA}$, $\mu = 0.69 \text{ cm}^{-1}$, $F(000) = 656$, $T = 294 \text{ K}$,
final $R = 0.072$ for 1037 observations.

After Lorenz corrections and polarization


corrections, the structure was solved using Direct Methods which make it possible to locate the majority of the nonhydrogen atoms of the molecule, the remaining atoms being located by Fourier differences and successive scaling operations. After isotropic refinement ($R = 0.125$) and then anisotropic refinement ($R = 0.095$), most of the hydrogen atoms are located using a Fourier difference (between 0.39 and 0.14 e\AA^{-3}), the others being positioned by calculation. The complete structure was refined by whole matrix (x, y, z, β_{ij} for C and O, x, y, z for H; 200 variables and 1037 observations; $w = 1/\sigma(F_o)^2 = [\sigma^2(I) + (0.04F_o^2)^2]^{-1/2}$) resulting in $R = 0.080$, $R_w = 0.072$ and $S_w = 1.521$ (residue $\Delta\rho \leq 0.21 \text{ e\AA}^{-3}$).

The scattering factors are taken from the International Tables of crystallography [International Tables for X-ray Crystallography (1974), Vol. IV, Birmingham: Kynoch Press (Current distributor D. Reidel, Dordrecht)]. The calculations were carried out on a Hewlett-Packard 9000-710 for the determination of the structure [Sheldrick, G.M. (1985), Crystallographic Computing 3: Data Collection, Structure Determination, Proteins and Databases, edited by G.M. Sheldrick, C. Krüger and R. Goddard, Oxford, Clarendon Press] and on a Digital MicroVax 3100 for the other calculations with the MOLEN suite of programs [Fair, C.K. (1990), MOLEN: An Interactive Intelligent System for Crystal Structure Analysis, Enraf-Nonius,

Delft, The Netherlands].

ORTEP DIAGRAM

[Johnson, C.K. (1965), ORTEP, Report ORNL-3794;
Oak Ridge National Laboratory, Tennessee, USA]



5


A (1S,2R,5S) - (+) -menthyl (2R,3R) -3-phenylglycidate sample, by treatment with sodium methoxide in methanol, made it possible to obtain the corresponding methyl phenylglycidate, the characteristics of which are as follows:

10

- $[\alpha]_D^{25} = +12$ (c = 1.15, chloroform)
- 400 MHz ^1H NMR (CDCl_3) (δ ppm): 7.40 (2H, d, J = 8 Hz), 7.32 (3H, m), 4.26 (1H, d, J = 4.6 Hz), 3.84 (1H, d, J = 4.6 Hz), 3.55 (3H, s).

Example 3:

(1S,2R,5S) - (+) -Menthyl (4S,5R) -2,4-diphenyl-
4,5-dihydrooxazole-5-carboxylate



15 mL (0.109 mol) of a 54% solution of
5 tetrafluoroboric acid in ether are added over 10 min to
a stirred solution, under an inert atmosphere at -65°C ,
of 30 g (0.0993 mol) of (1S,2R,5S) - (+) -menthyl (2R,3R) -
3-phenylglycidate and 305 mL (2.98 mol) of benzonitrile
in 1.5 L of anhydrous dichloromethane. The reaction is
10 allowed to continue at -65°C for 1 h and, after
monitoring by T.L.C., 300 mL of a saturated aqueous
sodium hydrogencarbonate solution are added and the
reaction mixture is allowed to return to room
temperature with stirring. After extracting the aqueous
15 phase with dichloromethane (2 x 200 mL), the combined
organic phases are washed with a saturated sodium
chloride solution (200 mL) and with water (50 mL) and
dried over MgSO_4 . After concentrating under reduced
pressure and removing the residual benzonitrile under
20 high vacuum at 50°C , the crude product obtained is
purified by silica gel chromatography (15-40 μm)

(eluent: cyclohexane/ethyl acetate, 20/1).


32 g of (1S,2R,5R)-(+)-menthyl (4S,5R)-2,4-diphenyl-4,5-dihydrooxazole-5-carboxylate are thus isolated in the form of a colourless syrup (Yd = 80%)

5 which exhibits the following characteristics:

- 400 MHz ^1H NMR (CDCl_3) (δ ppm): 8.10 (2H, d, $J = 7.1$ Hz), 7.54 (1H, t, $J = 7.4$ Hz), 7.46 (2H, t, $J = 7.4$ Hz), 7.34 (5H, m), 5.40 (1H, d, $J = 6.4$ Hz), 4.88 (1H, d, $J = 6.4$ Hz), 4.85 (1H, dt, $J = 10.9$ and 4.4 Hz), 2.09 (1H, m), 1.84 (1H, m, $J = 7$ and 2.7 Hz), 1.71 (1H, m), 1.69 (1H, m), 0.94 (3H, d, $J = 6.5$ Hz), 0.9 (1H, m), 0.85 (3H, d, $J = 7$ Hz), 0.77 (3H, d, $J = 7$ Hz).

Example 4:

15 (4S,5R)-2,4-Diphenyl-4,5-dihydrooxazole-5-carboxylic acid



25 mL of a solution of 6 g (43.2 mmol) of potassium carbonate in osmosed water are added to a stirred solution at room temperature of 3.5 g (8.64 mmol) of (1S,2R,5S)-(+)-menthyl (4S,5R)-2,4-diphenyl-4,5-dihydrooxazole-5-carboxylate in methanol (70 mL) and the reaction is left to continue for 16 h at room temperature. After monitoring by T.L.C., the


reaction mixture is concentrated under reduced pressure. The aqueous phase thus obtained is washed with dichloromethane (3 x 100 mL), acidified to pH 2 by slow addition of 20 mL of a 1M aqueous HCl solution and extracted with ethyl acetate (3 x 100 mL). The combined organic extraction phases are dried (MgSO₄) and concentrated under reduced pressure.

2.26 g of (4S,5R)-2,4-diphenyl-4,5-dihydrooxazole-5-carboxylic acid are thus obtained in the form of a white powder (Yd = 98%) which exhibits the following characteristics:

- $[\alpha]_D^{22} = +27.7$ (c = 0.99, CH₂Cl₂/MeOH, 1/1)
- F = 201-202°C
- 400 MHz ¹H NMR (d₆-DMSO) (δ ppm): 7.99 (2H, d, J = 7.3 Hz), 7.64 (1H, t, J = 7.4 Hz), 7.55 (2H, t, J = 7.7 Hz), 7.36 (5H, m), 5.40 (1H, d, J = 6.3 Hz), 4.99 (1H, d, J = 6.4 Hz).

Example 5:

(1S,2R,5S) - (+) -Menthyl (2R,3S) -N-benzoyl-3-phenylisoserinate



15 mL of a 1M aqueous HCl solution are added to a stirred solution at room temperature of 1 g (2.47 mmol) of (1S,2R,5S) - (+) -menthyl (4S,5R) -2,4-


diphenyl-4,5-dihydrooxazole-5-carboxylate in a mixture of methanol (15 mL) and tetrahydrofuran (15 mL). The reaction mixture is brought for 1 h to reflux and, after monitoring by T.L.C. and returning to room temperature, a saturated aqueous sodium hydrogencarbonate solution (45 mL) is gradually added until a basic pH is obtained. After stirring for 48 h at room temperature, the aqueous phase obtained after concentrating under reduced pressure is extracted with dichloromethane (100 mL). The aqueous phase is washed with a saturated sodium chloride solution (2 x 50 mL), dried over MgSO₄ and concentrated under reduced pressure and the residue obtained is chromatographed on silica gel (15-40 μ m) (eluent: dichloromethane/methanol, 95/05).

0.835 g of (1S,2R,5S)-(+)-menthyl (2R,3S)-N-benzoyl-3-phenylisoserinate is thus isolated in the form of a white solid (Yd = 80%) which exhibits the following characteristics:

• 400 MHz ¹H NMR (CDCl₃) (δ ppm): 7.77 (2H, d, J = 7.2 Hz), 7.51 (1H, t, J = 7.3 Hz), 7.45 (4H, m), 7.36 (2H, t, J = 7.2 Hz), 7.29 (1H, t, J = 7.2 Hz), 7.04 (1H, d, J = 9.2 Hz), 5.78 (1H, dd, J = 9.2 and 2.1 Hz), 4.79 (1H, dt, J = 10.9 and 4.4 Hz), 4.63 (1H, broad s), 3.35 (1H, broad s), 1.81 (2H, m), 1.67 (3H, m), 1.5 to 1.36 (2H, m), 1.09 to 0.91 (2H, m), 0.89 (3H, d, J = 6.9 Hz), 0.77 (3H, d, J = 6.5 Hz), 0.74 (3H, d, J = 6.9 Hz).

Example 6:

(1S,2R,5S) - (+) -Menthyl (2R,3S) -N-benzoyl-O-triethylsilyl-3-phenylisoserinate



0.255 g (2.08 mmol) of 4-dimethylamino-
 5 pyridine is added to a solution of 0.8 g (1.89 mmol) of
 (1S,2R,5S) - (+) -menthyl (2R,3S) -N-benzoyl-3-
 phenylisoserinate in 10 mL of anhydrous
 dichloromethane. After stirring for a few minutes at
 room temperature, 477 μ L (2.84 mmol) of triethylsilyl
 10 chloride are added over 5 min. After stirring for 1 h
 at room temperature and monitoring by T.L.C., the
 reaction mixture is diluted with 100 mL of
 dichloromethane. The organic phase is washed with a
 saturated aqueous sodium hydrogencarbonate solution
 15 (2 \times 20 mL) and with a saturated sodium chloride
 solution (50 mL), dried over MgSO_4 and concentrated
 under reduced pressure. After purifying the residue
 obtained by silica gel chromatography (15-40 μ m).
 (eluent: cyclohexane/ethyl acetate, 10/1), 0.74 g of
 20 (1S,2R,5S) - (+) -menthyl (2R,3S) -N-benzoyl-O-
 triethylsilyl-3-phenylisoserinate is obtained in the
 form of a colourless syrup (Yd = 75%).


The compound obtained exhibits the following

characteristics:

- 400 MHz ^1H NMR (CDCl_3) (δ ppm): 7.82 (2H, d, $J = 7$ Hz), 7.52 (1H, t, $J = 7.4$ Hz), 7.45 (2H, t, $J = 7$ Hz), 7.37 (2H, d, $J = 7.2$ Hz), 7.32 (2H, t, $J = 7.2$ Hz), 7.26 (2H, m), 5.60 (1H, dd), 4.73 (1H, dt, $J = 11$ and 4.3 Hz), 1.88 to 1.67 (m), 1.44 (2H, m), 1.06 and 0.87 (m), 0.80 (m), 0.67 (3H, d, $J = 7$ Hz), 0.62 to 0.34 (m).

Example 7:

(2R,3S)-N-Benzoyl-O-triethylsilyl-3-phenylisoserine



A solution of 0.644 g (4.655 mmol) of sodium carbonate in 10 mL of osmosed water is added to a stirred solution at room temperature of 0.5 g (0.931 mmol) of (1S,2R,5S)-(+)-menthyl (2R,3S)-N-benzoyl-O-triethylsilyl-3-phenylisoserinate in 15 mL of methanol. After stirring for 16 h at room temperature and monitoring by T.L.C., the reaction mixture is concentrated under reduced pressure and the residual aqueous phase is washed with dichloromethane (3 x 50 mL) and then acidified to pH 2 by slow addition of a 1M aqueous HCl solution (10 mL). The aqueous phase is extracted with ethyl acetate (3 x 50 mL) and the combined organic phases are dried over MgSO_4 and


concentrated under reduced pressure.

0.320 g of (2R,3S)-N-benzoyl-O-triethylsilyl-3-phenylisoserine is obtained in the form of a white powder (Yd = 90%) which exhibits the following characteristics:

- 400 MHz ^1H NMR (d_6 -DMSO) (δ ppm): 8.46 (1H, d, $J = 9.3$ Hz), 7.82 (2H, d, $J = 7.1$ Hz), 7.54 (1H, t, $J = 7.2$ Hz), 7.47 (4H, m), 7.32 (2H, t), 7.36 (1H, t), 5.44 (1H, dd, $J = 9.2$ and 5.5 Hz), 4.64 (1H, d, $J = 5.6$ Hz), 0.77 (9H, m), 0.45 (6H, m).

Example 8:

(1S,2R,5S) - (+) -Menthyl (2R,3S) -N-benzoyl-O-(2,2,2-trichloroethoxy)carbonyl-3-phenylisoserinate



480 mg (3.96 mmol) of 4-dimethylaminopyridine are added to a stirred solution at room temperature under an inert atmosphere of 1.38 g (3.3 mmol) of (1S,2R,5S) - (+) -menthyl (2R,3S) -N-benzoyl-3-phenylisoserinate in 30 mL of anhydrous dichloromethane. After stirring for 10 min, 540 μL (3.96 mmol) of 2,2,2-trichloroethoxycarbonyl chloride are added over 5 min. After stirring for 2 h at room temperature and monitoring by T.L.C., the organic phase is washed with a saturated sodium hydrogencarbonate.


solution (2 x 10 mL) and with a saturated sodium chloride solution (10 mL), dried over MgSO₄ and concentrated under reduced pressure. After purifying the residue obtained by silica gel chromatography (15-40 μm) (eluent: cyclohexane/ethyl acetate, 5/1), 1.60 g of (1S,2R,5S) - (+) -menthyl (2R,3S) - N-benzoyl-O-(2,2,2-trichloroethoxy)carbonyl-3-phenylisoserinate are obtained in the form of a colourless syrup (Yd = 82%).

The compound obtained exhibits the following characteristics:

- 400 MHz ¹H NMR (CDCl₃) (δ ppm): 7.82 (2H, d, J = 7.4 Hz), 7.53 (1H, t, J = 7.4 Hz), 7.44 (4H, m), 7.35 (2H, t, J = 7 Hz), 7.29 (1H, t, J = 7 Hz), 7.09 (1H, d, J = 9.3 Hz), 6.0 (1H, dd, J = 9.3 and 2.5 Hz), 5.45 (1H, d, J = 2.6 Hz), 4.78 and 4.72 (2H, 2d, J = 11.9 Hz), 4.77 (1H, m), 1.85 (1H, m), 1.79 (1H, m), 1.65 (2H, m), 1.43 (1H, m), 1.02 (1H, m), 0.96 (1H, m), 0.86 (1H, m), 0.83 (3H, d, J = 7 Hz), 0.78 (3H, d, J = 6.5 Hz), 0.68 (3H, d, J = 6.9 Hz).

Example 9:

(1S,2R,5S) - (+) -Menthyl (4S,5R) - 4-phenyloxazolidin-2-one-5-carboxylate




1 mL (7.28 mmol) of 1,8-diazabicyclo[5,4,0]-undec-7-ene is added to a stirred solution at room temperature under an inert atmosphere of 3.96 g (6.62 mmol) of (1S,2R,5S)-(+)-menthyl (2R,3S)-N-benzoyl-O-(2,2,2-trichloroethoxy) carbonyl-3-phenylisoserinate in 30 mL of anhydrous dichloromethane. After stirring for 30 min at room temperature, the organic phase is washed with 10 mL of a saturated sodium chloride solution, dried over MgSO₄ and concentrated under reduced pressure. After purifying the residue by silica gel chromatography (15-40 μ m) (eluent: cyclohexane/ethyl acetate, 7/3), 2.18 g of the compound cited in the title are obtained in the form of a yellow syrup (Yd = 95%).

The compound obtained exhibits the following characteristics:

- 400 MHz ¹H NMR (CDCl₃) (δ ppm): 7.40 (5H, m), 6.09 (1H, s), 4.93 (1H, d, J = 5.3 Hz), 4.86 (1H, dt, J = 11 and 4.4 Hz), 4.73 (1H, d, J = 5.4 Hz), 2.05 (1H, m), 1.81 (1H, m), 1.71 (2H, m), 1.54 to 1.41 (3H, m), 1.07 (2H, m), 0.94 (3H, d, J = 6.5 Hz), 0.88 (3H, d, J = 7 Hz), 0.77 (3H, d, J = 7 Hz).

Example 10:

(1S,2R,5S) - (+) -Menthyl (4S,5R) -N-t-butoxycarbonic-4-phenyloxazolidin-2-one-5-carboxylate



3.8 mL (6.07 mmol) of a 1.6M solution of
 5 n-butyllithium in hexane are added to a stirred
 solution at -40°C under an inert atmosphere of 1.91 g
 (5.52 mmol) of (1S,2R,5S) - (+) -menthyl (4S,5R) -4-
 phenyloxazolidin-3-one-5-carboxylate in 20 mL of
 anhydrous tetrahydrofuran. After stirring for 10 min at
 10 -40°C, a solution of 1.81 g (8.28 mmol) of
 t-butoxycarbonic anhydride in solution in 5 mL of
 tetrahydrofuran is added and the reaction mixture is
 allowed to return to room temperature over 15 min.
 After diluting with 50 mL of dichloromethane and
 15 washing with a 2% aqueous HCl solution until a pH = 5
 is obtained, the organic phase is dried (MgSO₄) and
 concentrated under reduced pressure. After purifying
 the crude product by silica gel chromatography
 (15-40 μm) (eluent: cyclohexane/ethyl acetate, 5/1),
 20 2.12 g of the compound cited in the title are obtained
 in the form of a colourless syrup (Yd = 86%).


The compound thus obtained exhibits the

following characteristics:

- 400 MHz ^1H NMR (CDCl_3) (δ ppm): 7.45 to 7.26 (5H, m), 5.19 (1H, d, $J = 3.7$ Hz), 4.86 (1H, dt, $J = 10.9$ and 4.5 Hz), 4.66 (1H, d, $J = 3.7$ Hz), 2.05 (1H, m), 1.79 (1H, m), 1.73 (2H, m), 1.62 to 1.24 (3H, m); 1.33 (9H, s), 1.11 (2H, m), 0.94 (3H, d, $J = 6.5$ Hz) and (1H, m), 0.89 (3H, d, $J = 7$ Hz), 0.77 (3H, d, $J = 7$ Hz).

Example 11:

(1S,2R,5S)-(+)-Menthyl (4S,5R)-3-N-benzoyl-4-phenyloxazolidin-3-one-5-carboxylate



0.25 mL (2.17 mmol) of benzoyl chloride is added to a stirred solution at room temperature under an inert atmosphere of 500 mg (1.45 mmol) of (1S,2R,5S)-(+)-menthyl (4S,5R)-4-phenyloxazolidin-3-one-5-carboxylate and 176 mg (1.16 mmol) of 4-pyrrolidinopyridine in 7 mL of anhydrous dichloromethane. After stirring for 3 h at 50°C, the reaction mixture is brought back to room temperature and diluted with 20 mL of dichloromethane. The organic phase is washed with 10 mL of a saturated sodium chloride solution, dried over MgSO_4 and concentrated under reduced pressure. After purifying the crude


product by silica gel chromatography (15-40 μ m)
(eluent: cyclohexane/ethyl acetate, 5/1), 300 mg of the
compound cited in the title are obtained in the form of
a colourless syrup (Yd = 46%).

5 The compound thus obtained exhibits the
following characteristics:

• 400 MHz ^1H NMR (CDCl_3) (δ ppm): 8.16 (2H, d,
J = 7.1 Hz), 7.68 (1H, t), 7.53 (4H, m), 7.43 (3H, m),
5.57 (1H, d, J = 4.4 Hz), 4.90 (1H, dt, J = 10.9 and
10 4.4 Hz), 4.85 (1H, d, J = 4.3 Hz), 2.07 (1H, m), 1.80
(1H, m), 1.72 (2H, m), 1.47 (3H, m), 1.09 (2H, m), 0.95
(3H, d, J = 6.5 Hz), 0.88 (3H, d, J = 7 Hz), 0.78 (3H,
d, J = 7 Hz).

Example 12:

15 (4S,5R)-3-N-Benzoyl-4-phenyloxazolidin-3-one-
5-carboxylic acid.



A solution of 75 mg (0.543 mmol) of potassium
carbonate in 1 mL of water is added to a stirred
mixture at room temperature of 120 mg (0.266 mmol) of
20 (1S,2R,5S)-(+)-menthyl (4S,5R)-3-N-benzoyl-4-
phenyloxazolidin-3-one-5-carboxylate in 2 mL of
methanol. After stirring for 30 min, the reaction
mixture is diluted with 10 mL of water and the aqueous


phase is washed with 5 mL of dichloromethane. After acidifying to pH = 4 by means of 1M HCl, the residual aqueous phase is extracted with ethyl acetate (3 x 10 mL). The combined organic phases are washed with 5 mL of a saturated sodium chloride solution, dried over MgSO₄ and concentrated under reduced pressure.

40 mg of (4S,5R)-3-N-benzoyl-4-phenyloxazolidin-3-one-5-carboxylic acid are obtained in the form of a white powder (Yd = 52%) which exhibits the following characteristics:

• 400 MHz ¹H NMR (d₆-DMSO) (δ ppm): 12.98 (1H, broad s), 7.95 (2H, d, J = 7.1 Hz), 7.63 (1H, t, J = 7.4 Hz), 7.50 (2H, t, J = 7.5 Hz), 7.42 (2H, m), 7.37 (3H, m), 4.90 (1H, d, J = 5 Hz), 4.77 (1H, d, J = 5 Hz).

Example 13:

(4S,5R)-4-Phenyloxazolidin-3-one-5-carboxylic acid



10 mL of a homogeneous solution of 360 mg (8.67 mmol) of NaOH, 3 mL of methanol and 0.5 mL of water in pyridine are rapidly added to a stirred solution at 0°C under an inert atmosphere of 300 mg (0.867 mmol) of (1S,2R,5S)-(+)-menthyl (4S,5R)-4-

phenyloxazolidin-2-one-5-carboxylate, 3 mL of methanol and then 0.5 mL of water in 6.5 mL of pyridine. After stirring for 20 min at 0°C, the reaction mixture is diluted with water (30 mL) and washed with dichloromethane (30 mL). After acidifying to a pH = 1, the residual aqueous phase is extracted with ethyl acetate (3 x 20 mL) and the combined organic phases are dried (MgSO₄) and concentrated under reduced pressure.


86 mg of (4S,5R)-4-phenyloxazolidin-3-one-5-carboxylic acid are thus obtained in the form of a yellow syrup (Yd = 53%) which exhibits the following characteristics:

- 400 MHz ¹H NMR (d₆-DMSO) (δ ppm): 13.33 (1H, broad s), 8.46 (1H, s), 7.38 (5H, m), 4.89 (1H, d, J = 5 Hz), 4.75 (1H, d, J = 5 Hz).

II. Baccatin III derivatives

Example 14:

7-O-Triethylsilyl-10-deacetylbaccatin III



6.2 mL (36.6 mmol) of triethylsilyl chloride are added over 10 min to a stirred solution, at room temperature and under an inert atmosphere, of 10 g (18.3 mmol) of 10-deacetylbaccatin III and 8.17 g

(54.9 mmol) of 4-pyrrolidinopyridine in 500 mL of anhydrous dichloromethane. After reacting for 3 h at room temperature, 10 g of crushed ice are added and the mixture is left stirring vigorously for 10 min. The residual organic phase is washed with water (200 mL), dried over MgSO_4 and concentrated under reduced pressure.


After treating the crude product obtained with the minimum amount of ethyl acetate, 11.2 g of 7-O-triethylsilyl-10-deacetylbaecatin III are obtained in the crystalline state (Yd = 92.3%).

The product thus obtained exhibits the following characteristics:

- 400 MHz ^1H NMR (CDCl_3) (δ ppm): 8.10 (2H, d, $J = 7.4$ Hz), 7.60 (1H, t, $J = 7.5$ Hz), 7.47 (2H, t, $J = 7.6$ Hz), 5.60 (1H, d, $J = 7$ Hz), 5.17 (1H, d, $J = 1.9$ Hz), 4.96 (1H, d, $J = 8$ Hz), 4.86 (1H, m), 4.41 (1H, dd, $J = 10.6$ and 6.6 Hz), 4.31 and 4.16 (2H, 2d, $J = 8.4$ Hz), 4.26 (1H, d, $J = 1.9$ Hz), 3.95 (1H, d, $J = 6.9$ Hz), 2.48 (1H, ddd, $J = 14.5$, 9.7 and 6.7 Hz), 2.29 (3H, s), 2.27 (2H, m), 2.08 (3H, s), 1.90 (1H, m), 1.73 (3H, s), 1.62 (1H, s), 1.08 (6H, s), 0.94 (9H, t, $J = 8$ Hz), 0.56 (6H, m).

Example 15:

7-O-Triethylgermanyl-10-deacetylbaccatin III




80 μ L (0.476 mmol) of triethylgermanyl chloride are added over 10 min to a stirred solution, at room temperature and under an inert atmosphere, of 100 mg (0.183 mmol) of 10-deacetylbaccatin III and 41 mg (0.275 mmol) of 4-pyrrolidinopyridine in 4 ml of anhydrous dichloromethane and the mixture is stirred at 50°C for 13 h. After cooling the reaction mixture and diluting with 15 mL of dichloromethane, 1 g of crushed ice is added and the mixture is left stirring vigorously for 10 min. The residual organic phase is washed with a saturated sodium hydrogencarbonate solution (5 mL) and a saturated sodium chloride solution (5 mL), dried over MgSO_4 and concentrated under reduced pressure. After chromatographing the crude product on silica gel (15-40 μ m) (eluent: cyclohexane/ethyl acetate, 25/75), 67 mg of 7-O-triethylgermanyl-10-deacetylbaccatin III are obtained in the form of a colourless syrup.

The product thus obtained exhibits the following characteristics:

• 400 MHz ^1H NMR (CDCl_3) (δ ppm): 8.09 (2H, d, $J = 7.1$ Hz), 7.60 (1H, t, $J = 7.4$ Hz), 7.48 (2H, t, $J = 7.6$ Hz), 5.63 (1H, d, $J = 7.1$ Hz), 5.24 (1H, s), 4.99 (1H, d, $J = 8$ Hz), 4.78 (1H, t), 4.32 (1H, d, $J = 8.3$), 4.28 (1H, m), 4.17 (2H, m), 3.97 (1H, d, $J = 7$ Hz), 2.59 (1H, m), 2.30 (3H, s), 2.24 (1H, m), 2.10 (1H, m), 2.03 (3H, s), 1.82 (1H, m), 1.73 (3H, s), 1.11 (9H, m), 1.0 (6H, t, $J = 7.7$ Hz).

Example 16:

10 7-O-(2,2,2-Trichloro-*t*-butoxycarbonyl)-10-deacetylbaccatin III



3.3 g (13.8 mmol) of 2,2,2-trichloro-*t*-butoxycarbonyl chloride are added over 2 h to a stirred solution at 40°C under an inert atmosphere of 5 g (9.19 mmol) of 10-deacetylbaccatin III and 1.1 mL of anhydrous pyridine in 250 mL of dry dichloromethane. After reacting for an additional 30 min and returning to room temperature, the organic solution is washed with a 2% aqueous HCl solution (30 mL), washed with osmosed water (2 x 100 mL), dried over MgSO_4 and concentrated under reduced pressure (Yd = 55%). After chromatographing the crude product on silica gel




5

10

Example 17:

a) 7-O-Triethylsilylbaccatin III



20


dichloromethane. After reacting for 2 h at room temperature and monitoring by T.L.C., 1 g of crushed ice is added and the mixture is left stirring vigorously for 10 min. The residual organic phase is washed with water (2 x 10 mL), dried over MgSO₄ and concentrated under reduced pressure. After silica gel chromatography (15-40 μ m) (eluent: cyclohexane/ethyl acetate, 60/40), 0.756 g of 7-O-triethylsilylbaccatin III is obtained in the form of a white powder (Yd = 70%).

The compound obtained exhibits the following characteristics:

- 400 MHz ¹H NMR (CDCl₃) (δ ppm): 8.11 (2H, d, J = 7.1 Hz), 7.6 (1H, t, J = 7.4 Hz), 7.48 (2H, t, J = 7.7 Hz), 6.46 (1H, s), 5.63 (1H, d, J = 7 Hz), 4.96 (1H, d, J = 8.1 Hz), 4.83 (1H, m), 4.49 (1H, dd, J = 10.4 and 6.7 Hz), 4.31 and 4.15 (2H, 2d, J = 8.3 Hz), 3.88 (1H, d, J = 7 Hz), 2.53 (1H, m), 2.29 (3H, s), 2.27 (2H, m), 2.19 (3H, d, J = 0.8 Hz), 2.18 (3H, s), 2.12 (1H, d), 1.88 (1H, m), 1.68 (3H, s), 1.65 (1H, s), 1.2 (3H, s), 1.04 (3H, s), 0.92 (9H, t), 0.59 (6H, m).

Example 18:

7-O-(2,2,2-Trichloro-t-butoxycarbonyl)-
baccatin III



50 μ L (0.695 mmol) of acetyl chloride are
5 added to a stirred solution at room temperature under
an inert atmosphere of 260 mg of 7-O-(2,2,2-trichloro-
t-butoxycarbonyl-10-deacetylbaccatin III and 127.5 mg
(1.04 mmol) of 4-dimethylaminopyridine in 2.5 mL of dry
dichloromethane. After reacting for 1 h at room
10 temperature, the organic phase is washed with a 2%
aqueous HCl solution until a pH = 6 is obtained, dried
over MgSO_4 and concentrated under reduced pressure.
After chromatographing the residue obtained on silica
gel (15-40 μ m) (eluent: cyclohexane/ethyl acetate,
15 6/4), 0.23 g of 7-O-(2,2,2-trichloro-t-
butoxycarbonyl)baccatin III is obtained in the solid
state (Yd = 83%).


The compound obtained exhibits the following
characteristics:

20 • 400 MHz ^1H NMR (CDCl_3) (δ ppm): 8.11 (2H, d,
 J = 7.1 Hz), 7.62 (1H, t, J = 7.4 Hz), 7.49 (2H, t,
 J = 7.6 Hz), 6.39 (1H, s), 5.64 (1H, d, J = 6.9 Hz),

5.61 (1H, dd, $J = 10.7$ and 7.2 Hz), 4.99 (1H, d, $J = 8.2$ Hz), 4.87 (1H, m), 4.33 and 4.16 (2H, 2d, $J = 8.4$ Hz), 4.02 (1H, d, $J = 6.9$ Hz), 2.64 (1H, ddd, $J = 14.4, 9.5$ and 7.2 Hz), 2.30 (3H, s) and (2H, m),
 5 2.17 (3H, s), 2.13 (3H, d, $J = 0.8$ Hz), 2.04 (1H, m),
 1.83 (3H, s), 1.63 (1H, s), 1.14 (3H, s), 1.09 (3H, s).

Example 19:

7-O-Phenoxyacetyl-10-deacetylbaccatin III



1.05 mL (7.5 mmol) of phenoxyacetyl chloride
 10 are added over 10 min to a stirred solution, at room
 temperature and under an inert atmosphere, of 1.03 g
 (1.88 mmol) of 10-deacetylbaccatin III and 0.6 mL
 (7.5 mmol) of anhydrous pyridine in 100 mL of dry
 dichloromethane. After reacting for 30 min at room
 15 temperature and monitoring by T.L.C., the organic
 solution is washed with a 2% aqueous HCl solution until
 a pH = 2 is obtained, washed with osmosed water
 (2 x 50 mL), dried over MgSO_4 and concentrated under
 reduced pressure (Yd = 70.5%). After chromatographing
 20 the crude product on silica gel (15-40 μm) (eluent:
 cyclohexane/ethyl acetate, 60/40), 7-O-phenoxyacetyl-
 10-deacetylbaccatin III is obtained in the form of a

white powder.


The product obtained exhibits the following characteristics:

- 400 MHz ^1H NMR (CDCl_3) (δ ppm): 8.09 (2H, d, $J = 7.3$ Hz), 7.61 (1H, t, $J = 7.4$ Hz), 7.48 (2H, t, $J = 7.6$ Hz), 7.31 (2H, t, $J = 7.7$ Hz), 6.99 (3H, m), 6.42 (1H, s), 5.61 (1H, d, $J = 7$ Hz), 4.97 (1H, d, $J = 7.8$ Hz), 4.86 (3H, m), 4.44 (1H, dd, $J = 10.6$ and 6.8 Hz), 4.30 and 4.15 (2H, 2d, $J = 8.4$ Hz), 3.86 (1H, d, $J = 7$ Hz), 2.56 (1H, m), 2.27 (3H, s), 2.27 (2H, m), 2.05 ((3H, s), 1.86 (1H, m), 1.68 (3H; s), 1.01 (3H, s), 0.98 (3H, s).

Example 20:

7,10-O-Di(phenoxycetyl)-10-deacetylbaccatin

15 III



0.5 mL (3.68 mmol) of phenoxycetyl chloride is added over 10 min to a stirred solution, at room temperature and under an inert atmosphere, of 500 mg (0.92 mmol) of 10-deacetylbaccatin III and 0.6 mL (7.36 mmol) of anhydrous pyridine in 50 mL of dry dichloromethane. After reacting for 6 h at room temperature and monitoring by T.L.C., the solution is

washed with a 2% aqueous HCl solution until a pH = 2 is obtained, washed with osmosed water (2 x 20 mL), dried over MgSO₄ and concentrated under reduced pressure.


After chromatographing the crude product on silica gel (15-40 μ m) (eluent: cyclohexane/ethyl acetate, 6/4), 0.55 g of 7-10-O-bis(phenoxyacetyl)-10-deacetylbaccatin III is obtained in the form of a white powder (Yd = 74%).

The product obtained exhibits the following characteristics:

- 400 MHz ¹H NMR (CDCl₃) (δ ppm): 8.09 (2H, d, J = 7.1 Hz), 7.61 (1H, t, J = 7.4 Hz), 7.48 (2H, t, J = 7.6 Hz), 7.29 (2H, t, J = 6.8 Hz), 7.22 (2H, t, J = 7.5 Hz), 6.96 (4H, m), 6.84 (2H, d, J = 7.9 Hz), 6.42 (1H, s), 5.69 (1H, dd, J = 10.5 and 7.1 Hz), 5.60 (1H, d, J = 6.9 Hz), 4.96 (1H, d, J = 8.2 Hz), 4.84 (1H, t, J = 7.4 Hz), 4.8 (2H, s), 4.65 and 4.41 (2H, 2d, J = 15.8 Hz), 4.32 and 4.14 (2H, 2d, J = 8.4 Hz), 3.98 (1H, d, J = 6.8 Hz), 2.65 (1H, m), 2.28 (3H, s), 2.26 (2H, m), 2.09 (3H, s), 1.80 (3H, s) and (1H, m), 0.98 (6H, s).

Example 21:

7-O-Phenoxyacetylbaccatin III



0.233 mL (3.27 mmol) of acetyl chloride is added over 10 min to a stirred solution, at room temperature and under an inert atmosphere, of 1.11 g (1.64 mmol) of 7-O-phenoxyacetyl-10-deacetylbaccatin III in 40 mL of anhydrous pyridine. After reacting for 16 h at room temperature and monitoring by T.L.C., the reaction mixture is diluted with 50 mL of osmosed water and the aqueous phase is extracted with ethyl acetate (3 x 30 mL). The combined organic phases are washed with water (2 x 20 mL), dried over MgSO_4 and concentrated under reduced pressure (Yd = 84.5%). After silica gel chromatography (15-40 μm) (eluent: cyclohexane/ethyl acetate, 60/40), 7-O-phenoxyacetyl-
baccatin III is obtained in the crystalline state.


The product obtained exhibits the following characteristics:

- 400 MHz ^1H NMR (CDCl_3) (δ ppm): 8.10 (2H, d, J = 7.1 Hz), 7.61 (1H, t, J = 7.4 Hz), 7.48 (2H, t, J = 7.7 Hz), 7.27 (2H, t, J = 8 Hz), 6.95 (3H, m), 6.26 (1H, s), 5.71 (1H, dd, J = 10.4 and 7.2 Hz), 5.62 (1H,

d, $J = 6.9$ Hz), 4.96 (1H, d, $J = 8.3$ Hz), 4.80 (1H, m),
 4.81 and 4.53 (2H, 2d, $J = 16$ Hz), 4.32 and 4.14 (2H,
 2d, $J = 8.5$ Hz), 4.0 (1H, d, $J = 6.9$ Hz), 2.64 (1H, m),
 2.29 (2H, m), 2.28 (3H, s), 2.24 (1H, d, $J = 5$ Hz),
 5. 2.16 (3H, s), 2.09 (3H, d, $J = 0.7$ Hz), 1.81 (1H, m),
 1.78 (3H, s), 1.13 (3H, s), 1.08 (3H, s).

Example 22;

7-10-0-(1,1,3,3-Tetraisopropyl-1,3-
 disiloxanediyl)-10-deacetylbaccatin III



10 1.28 ml (2.05 mmol) of *n*-butyllithium as a
 1.6M solution in hexane are added over 10 min to a
 stirred solution, at -40°C and under an inert
 atmosphere, of 500 mg (0.93 mmol) of 10-deacetyl-
 baccatin III in 20 mL of anhydrous tetrahydrofuran.
 15 After stirring for 5 min, 350 μL (1.12 mmol) of 1,3-
 dichloro-1,1,3,3-tetraisopropyldisiloxane are added and
 the reaction mixture is allowed to return to room
 temperature over 20 min. After stirring for 1 h at room
 temperature, 225 mg (2.05 mmol) of 4-dimethylamino-
 20 pyridine are added and the reaction mixture is left


stirring for an additional 1 h. After adding 20 mL of a saturated aqueous sodium chloride solution, the mixture is extracted with dichloromethane (3 x 30 mL). The combined organic phases are washed with a saturated aqueous sodium chloride solution (20 ml), dried over MgSO₄ and concentrated under reduced pressure. After purifying by silica gel chromatography (15-40 μ m) (eluent: cyclohexane/ethyl acetate, 60/40), 480 mg of 7,10-O-(1,1,3,3-tetraisopropyl-1,3-disyloxanediyl)-10-deacetylbaecatin III are obtained in the amorphous state (Yd = 65%).

The product obtained exhibits the following characteristics:

- 400 MHz ¹H NMR (CDCl₃) (δ ppm): 8.10 (2H, d, J = 7.2 Hz), 7.60 (1H, t, J = 7.4 Hz), 7.47 (2H, t, J = 7.6 Hz), 5.60 (1H, s), 5.59 (1H, d), 4.97 (1H, d, J = 7.9 Hz), 4.87 (1H, m), 4.68 (1H, dd, J = 10.4 and 6.9 Hz), 4.30 and 4.17 (2H, 2d, J = 8.5 Hz), 3.92 (1H, d, J = 7.1 Hz), 2.49 (1H, m), 2.28 (3H, s), 2.27 (1H, m), 2.04 (1H, m), 1.91 (1H, m), 1.67 (3H, s), 1.55 (1H, s), 1.32 to 0.85 (34H, m).

Example 23:

13-O-[[(4S,5R)-2,4-Diphenyl-4,5-dihydroxazol-5-yl]carbonyl]-7-O-triethylsilylbaccatin III



2.06 g (10 mmol) of dicyclohexylcarbodiimide
 5 are added to a stirred solution, at room temperature
 and under an inert atmosphere, of 2.67 g (10 mmol) of
 (4S,5R)-2,4-diphenyl-4,5-dihydroxazol-5-carboxylic acid
 in 55 mL of anhydrous toluene. After stirring for
 5 min, 3.5 g (5 mmol) of 7-O-triethylsilylbaccatin III
 10 and 0.61 g (5 mmol) of 4-dimethylaminopyridine are
 added and the reaction mixture is brought to 70°C for
 1 h. After returning to room temperature and removing
 the insoluble materials by filtration, the organic
 phase is concentrated under reduced pressure. After
 15 purifying the crude product by silica gel
 chromatography (15-25 μ m) (eluent: cyclohexane/ethyl
 acetate, 90/10).


4.62 g of 13-O-[[(4S,5R)-2,4-diphenyl-4,5-
 dihydroxazol-5-yl]carbonyl]-7-O-triethylsilylbaccatin
 20 III are obtained in the crystalline state (Yd = 97%).

The compound thus obtained exhibits the following characteristics:

• 400 MHz ^1H NMR (CDCl_3) (δ , ppm): 8.23 (2H, d, $J = 7.2$ Hz), 8.07 (2H, d, $J = 7.3$ Hz), 7.63 (1H, t, $J = 7.4$ Hz), 7.58 (1H, t, $J = 7.4$ Hz), 7.49 (4H, m), 7.38 (5H, m), 6.42 (1H, s), 6.18 (1H, t, $J = 8.2$ Hz), 5.68 (1H, d, $J = 7.1$ Hz), 5.60 (1H, d, $J = 6.5$ Hz), 4.95 (2H, d), 4.50 (1H, dd, $J = 10.5$ and 6.7 Hz), 4.29 (1H, d, $J = 8.4$ Hz), 4.14 (1H, d, $J = 8.4$ Hz), 3.83 (1H, d, $J = 7.1$ Hz), 2.55 (1H, m), 2.37 (1H, dd, $J = 15.3$ and 9.3 Hz), 2.26 (1H, dd, $J = 15.3$ and 8.6 Hz), 2.16 (3H, s), 2.07 (3H, s), 1.99 (3H, s), 1.89 (1H, m), 1.72 (1H, s), 1.69 (3H, s), 1.23 (3H, s), 1.19 (3H, s), 0.92 (9H, t, $J = 8$ Hz), 0.57 (6H, m).

Example 24:

13-O-[[4S,5R)-2,4-Diphenyl-4,5-dihydrooxazol-5-yl]carbonyl]-7-O-phenoxycetylbaccatin III



380 mg (1.84 mmol) of dicyclohexylcarbodiimide are added to a stirred solution, at room temperature and under an inert atmosphere, of 490 mg

(1.83 mmol) of (4S,5R)-2,4-diphenyl-4,5-dihydrooxazol-5-carboxylic acid in 10 mL of anhydrous toluene. After stirring for 5 min, 660 mg (0.92 mmol) of 7-O-phenoxyacetylbaecatin III and 112 mg (0.92 mmol) of 4-dimethylaminopyridine are added and the reaction mixture is brought to 70°C for 2 h.


After returning to room temperature and removing the insoluble materials by filtration, the organic phase is concentrated under reduced pressure. After purifying the crude product by silica gel chromatography (15-40 μ m) (eluent: cyclohexane/ethyl acetate, 99/1), 800 mg of 13-O-[(4S,5R)-2,4-diphenyl-4,5-dihydrooxazol-5-yl]carbonyl]-7-O-phenoxyacetylbaecatin III are obtained in the crystalline state (Yd = 90%).

The compound thus obtained exhibits the following characteristics:

- 400 MHz ^1H NMR (CDCl_3) (δ ppm): 8.18 (2H, d, $J = 7$ Hz), 8.07 (2H, d, $J = 7.3$ Hz), 7.63 (1H, t, $J = 7.4$ Hz), 7.59-7.32 (10H, m), 7.28 (2H, t, $J = 7.5$ Hz), 6.94 (3H, m), 6.23 (1H, s) and (1H, m), 5.70 (1H, dd, $J = 10.4$ and 7.1 Hz), 5.67 (1H, d, $J = 7.3$ Hz), 5.58 (1H, d, $J = 7$ Hz), 4.93 (2H, d), 4.79 and 4.53 (2H, 2d, $J = 15.9$ Hz), 4.30 and 4.13 (2H, 2d, $J = 8.5$ Hz), 3.97 (1H, d, $J = 6.9$ Hz), 2.67 (1H, m), 2.38 (1H, dd, $J = 15.2$ and 9.3 Hz), 2.26 (1H, dd, $J = 15.2$ and 8.4 Hz), 2.15 (3H, s), 2.02 (3H, s), 1.95 (3H, s) and (1H, m), 1.80 (3H, s), 1.74 (1H, s), 1.25 (3H, s), 1.17 (3H, s).

Example 25:

13-O-[[(4S,5R)-2,4-Diphenyl-4,5-dihydrooxazol-5-yl]carbonyl]-7-O-(2,2,2-trichloro-t-butoxycarbonyl)baccatin III



27 mg (0.13 mmol) of dicyclohexylcarbodiimide are added to a stirred solution, at room temperature and under an inert atmosphere, of 35 mg of (4S,5R)-2,4-diphenyl-4,5-dihydrooxazol-5-carboxylic acid in 3 mL of anhydrous toluene. After stirring for 5 min, 51 mg (0.065 mmol) of 7-O-(2,2,2-trichloro-t-butoxycarbonyl)baccatin III and 8 mg (0.065 mmol) of 4-dimethylaminopyridine are added and the mixture is brought to 70°C for 1 h. After returning to room temperature and removing the insoluble materials by filtration, the organic phase is concentrated under reduced pressure and the residue obtained is purified by silica gel chromatography (15-40 μ m) (eluent: cyclohexane/ethyl acetate, 9/1).


0.99 g of the compound cited in the title is thus obtained in the form of a white solid (Yd = 67%)

which exhibits the following characteristics:

• 400 MHz ^1H NMR (CDCl_3) (δ ppm): 8.18 (2H, d, $J = 7.2$ Hz), 8.07 (2H, d, $J = 7.3$ Hz), 7.65 (1H, t, $J = 7.4$ Hz), 7.59 (1H, t, $J = 7.3$ Hz), 7.52 (4H, m), 7.39 (5H, m), 6.35 (1H, s), 6.24 (1H, t, $J = 8.4$ Hz), 5.68 (1H, d, $J = 7.1$ Hz), 5.59 (1H, d, $J = 7$ Hz) and (1H, dd), 4.95 (1H, d), 4.94 (1H, d, $J = 7$ Hz), 4.31 and 4.15 (2H, 2d, $J = 8.4$ Hz), 3.97 (1H, d, $J = 6.9$ Hz), 2.64 (1H, m), 2.37 (1H, dd, $J = 15.1$ and 6 Hz), 2.27 (1H, dd, $J = 15.2$ and 8.5 Hz), 2.16 (3H, s), 2.01 (3H, s), 1.98 (3H, s), 1.83 (3H, s), 1.72 (1H, s), 1.25 (3H, s), 1.18 (3H, s).

Example 26:

13-*O*-[[4*S*,5*R*)-2,4-Diphenyl-4,5-dihydrooxazol-5-yl]carbonyl]-7,10-*O*-(1,1,3,3-tetraisopropyl-1,3-disiloxanediy)l)-10-deacetylbaecatin III



7 mg (0.06 mmol) of dicyclohexylcarbodiimide are added to a stirred solution, at room temperature and under an inert atmosphere, of 4 mg (0.015 mmol) of (4*S*,5*R*)-2,4-diphenyl-4,5-dihydrooxazol-5-carboxylic


acid in 0.5 mL of anhydrous toluene. After stirring for 5 min, a solution of 5 mg (0.0065 mmol) of 7,10-0-(1,1,3,3-tetraisopropyl-1,3-disiloxanediyl)-10-deacetylbaccatin III and of 1 mg (0.0078 mmol) of 4-dimethylaminopyridine in 1 mL of anhydrous toluene is added. After stirring for 20 min at room temperature, the mixture is brought to 50°C for an additional 20 min. After returning to room temperature, the organic phase is diluted with 5 mL of dichloromethane, washed with 2 mL of a saturated aqueous sodium chloride solution, dried over MgSO₄ and concentrated under reduced pressure. After purifying the crude product by silica gel chromatography (15-25 μ m) (eluent: cyclohexane/ethyl acetate, 7/3), 6 mg of the derivative cited in the title are obtained (Yd = 90%) in the amorphous state.

The compound obtained exhibits the following characteristics:

- 400 MHz ¹H NMR (CDCl₃) (δ ppm): 8.21 (2H, d, J = 7.2 Hz), 8.07 (2H, d, J = 7.6 Hz), 7.63 (1H, t, J = 7.5 Hz), 7.59 (1H, t, J = 7.4 Hz), 7.50 (2H, t, J = 7.4 Hz), 7.39 (5H, m), 6.26 (1H, t), 5.64 (1H, d, J = 7 Hz), 5.59 (1H, d, J = 6.9 Hz), 5.54 (1H, s), 4.93 (1H, d, J = 6.8 Hz) and (1H, m), 4.68 (1H, dd), 4.28 and 4.16 (2H, 2d, J = 8 Hz), 3.84 (1H, d, J = 7.3 Hz), 2.48 (1H, m), 2.35 and 2.25 (2H, 2dd), 2.02 (3H, s), 1.88 (3H, s) and (1H, m), 1.67 (3H, s), 1.63 (1H, s), 1.30 to 0.90 (34H, m).

Example 27:

13-O-[[[(4S,5R)-3-N-Benzoyl-4-phenyloxazolidin-3-one-5-yl]carbonyl]-7-O-triethylsilylbaccatin III




28 mg (0.136 mmol) of dicyclohexylcarbodiimide are added to a stirred solution at room temperature under an inert atmosphere of 40 mg (0.137 mmol) of (4S,5R)-3-N-benzoyl-4-phenyloxazolidin-3-one-5-carboxylic acid in 2 mL of anhydrous toluene. After stirring for 5 min, 30 mg (0.043 mmol) of 7-O-triethylsilylbaccatin III and 8 mg (0.066 mmol) of 4-dimethylaminopyridine are added and the reaction mixture is brought to 60°C for 13 h. After returning to room temperature, the reaction mixture is diluted with 10 mL of dichloromethane and the organic phase is washed with 5 mL of a saturated sodium chloride solution, dried over MgSO₄ and concentrated under reduced pressure. After purifying by silica gel chromatography (15-14 μm) (eluent: cyclohexane/ethyl acetate, 2/1), 13 mg of the derivative cited in the title are obtained in the amorphous state (Yd = 31%).

The compound obtained exhibits the following characteristics:

- 400 MHz ^1H NMR (CDCl_3) (δ ppm): 8.06 (2H, d, $J = 7.3$ Hz), 7.72 (2H, d, $J = 7$ Hz), 7.63 (1H, t, $J = 7.4$ Hz), 7.58 (1H, t, $J = 7.4$ Hz), 7.54 to 7.44 (8H, m), 7.40 (1H, t), 6.44 (1H, s), 6.33 (1H, t), 5.73 (1H, d, $J = 5.7$ Hz), 5.67 (1H, d, $J = 5.7$ Hz), 4.96 (1H, d, $J = 5.8$ Hz), 4.88 (1H, d, $J = 8.3$ Hz), 4.45 (1H, dd, $J = 10.4$ and 6.6 Hz), 4.27 and 4.12 (2H, 2d, $J = 8.3$ Hz), 3.80 (1H, d, $J = 7$ Hz), 2.50 (1H, m), 2.26 (2H, m), 2.19 (3H, s), 2.07 (3H, s), 1.98 (3H, s), 1.85 (1H, m), 1.76 (1H, s), 1.67 (3H, s), 1.24 (3H, s), 1.23 (3H, s), 0.91 (9H, t, $J = 7.9$ Hz), 0.56 (6H, m).

Example 28:

- 15 13-O-[[(4S,5R)-4-Phenyloxazolidin-3-one-5-yl]carbonyl]-7,10-O-di(phenoxycetyl)-10-deacetylbaaccatin III



- 20 65 mg (0.315 mmol) of dicyclohexylcarbodiimide are added to a stirred solution, at room temperature and under an inert atmosphere, of 78 mg

(0.293 mmol) of (4S,5R)-2,4-diphenyl-4,5-dihydrooxazol-5-carboxylic acid in 3 mL of anhydrous toluene. After stirring for 5 min, a solution of 237 mg (0.293 mmol) of 7,10-O-bis(phenoxyacetyl)-10-deacetylbaecatin III and 36 mg (0.295 mmol) of 4-dimethylaminopyridine in 3 mL of toluene is added and the reaction mixture is brought to 60°C for 1 h. After returning to room temperature and removing the insoluble materials by filtration, the organic phase is concentrated under reduced pressure and the crude product obtained is purified by silica gel chromatography (15-40 μ m) (eluent: cyclohexane/ethyl acetate, 1/1).

280 mg of the compound cited in the title are thus obtained in the amorphous state (Yd = 90%), which compound exhibits the following characteristics:

- 400 MHz ^1H NMR (CDCl_3) (δ ppm): 8.18 (2H, d, $J = 7$ Hz), 8.06 (2H, d, $J = 7.1$ Hz), 7.64 (1H, t, $J = 7.4$ Hz), 7.58 (1H, t, $J = 7.3$ Hz), 7.51 (4H, m), 7.39 (5H, m), 7.25 (4H, m), 6.96 (4H, m), 6.85 (2H, d, $J = 8$ Hz), 6.33 (1H, s), 6.19 (1H, t, $J = 9$ Hz), 5.68 (1H, dd, $J = 10.5$ and 7.1 Hz), 5.65 (1H, d, $J = 6.9$ Hz), 5.59 (1H, d, $J = 7$ Hz), 4.93 (2H, d, $J = 7.1$ Hz), 4.79 (2H, s), 4.63 and 4.40 (2H, 2d, $J = 15.9$ Hz), 4.30 and 4.13 (2H, 2d, $J = 8.4$ Hz), 3.94 (1H, d, $J = 6.9$ Hz), 2.68 (1H, m), 2.37 (1H, dd, $J = 15.3$ and 9.3 Hz), 2.24 (1H, dd, $J = 15.3$ and 8.7 Hz), 2.02 (3H, s), 1.95 (3H, s), 1.80 (3H, s) and (1H, m), 1.69 (1H, s), 1.12 (3H, s), 1.01 (3H, s).

III. Hemisynthesis

Example 29:

Preparation of paclitaxel

a) From 13-O-[[[(4S,5R)-2,4-diphenyl-4,5-dihydrooxazol-5-yl]carbonyl]-7-O-triethylsilylbaccatin
5 III

0.6 L (0.6 mol) of a 1M aqueous HCl solution is added to a stirred solution, at room temperature and under an inert atmosphere, of 90 g (0.095 mol) of 13-O-
10 [[[(4S,5R)-2,4-diphenyl-4,5-dihydrooxazol-5-yl]-carbonyl]-7-O-triethylsilylbaccatin III in a mixture of tetrahydrofuran (1.2 L) and methanol (1.2 L) and the reaction mixture is stirred at room temperature for 4 h 30. After adding 3.5 L of a saturated aqueous
15 sodium hydrogencarbonate solution, the solution is kept homogeneous by addition of 6 L of tetrahydrofuran and 6 L of water and the reaction mixture is stirred for an additional 1 h 30. After adding 15 L of ethyl acetate and 15 L of osmosed water, the residual aqueous phase
20 is extracted with ethyl acetate (15 L). The organic phase is dried over MgSO₄ and concentrated under reduced pressure and the crude product thus obtained is purified by silica gel chromatography (15-40 μm) (eluent: cyclohexane/ethyl acetate, 1/1).

25 75 g of taxol are thus isolated in the crystalline state (Yd = 95%), the characteristics of which are in every respect in accordance with the literature data.

b) From 13-O-[[[(4S,5R)-2,4-diphenyl-4,5-dihydrooxazol-5-yl]carbonyl]-7-O-(2,2,2-trichloro-t-butoxycarbonyl)baccatin III.

90 μ L (0.09 mmol) of a 1M aqueous HCl solution are added to a stirred solution, at room temperature and under an inert atmosphere, of 15 mg (0.0148 mmol) of 13-O-[[[(4S,5R)-2,4-diphenyl-4,5-dihydrooxazol-5-yl]carbonyl]-7-O-(2,2,2-trichloro-t-butoxycarbonyl)baccatin III in a mixture of tetrahydrofuran (0.18 mL) and methanol (0.18 mL) and the reaction mixture is stirred at room temperature for 8 h. After adding 0.6 mL of a saturated aqueous sodium hydrogencarbonate solution, the solution is kept homogeneous by addition of 1 mL of tetrahydrofuran and 1 mL of water and the reaction mixture is stirred for an additional 1 h 30. After adding 2.5 mL of ethyl acetate and 2.5 mL of osmosed water, the residual aqueous phase is extracted with ethyl acetate (2.5 mL). The combined organic phases are dried over MgSO_4 and concentrated under reduced pressure.

14 mg of 7-O-(2,2,2-trichloro-t-butoxycarbonyl)taxol are thus obtained in the crude state ($\text{Yd} = 93\%$), which product is used without additional purification in the following stage.

30 μ L (0.525 mmol) of acetic acid and 22.5 mg (0.344 mmol) of zinc powder are added to a stirred solution at room temperature of 13 mg (0.0128 mmol) of 7-O-(2,2,2-trichloro-t-butoxycarbonyl)taxol in 2 mL of


ethyl acetate. After stirring for 2 h 30 at room temperature and monitoring by T.L.C., and after diluting the reaction mixture with 3 mL of ethyl acetate, the organic phase is washed with osmosed water (1 mL), with a saturated aqueous sodium hydrogencarbonate solution (1 mL) and again with water, dried over MgSO_4 and concentrated under reduced pressure.

After chromatographing the crude product on silica gel (15-40 μm) (eluent: cyclohexane/ethyl acetate, 6/4), 9.5 mg of taxol are thus isolated in the crystalline state ($\text{Yd} = 89\%$).

Material which is outside the scope of the claims does not constitute a part of the claimed invention.

CLAIMS

1. Process for the preparation of taxane side chain precursors in which a cis- β -arylglycidate derivative of general formula I



5 in which

Ar represents an aryl radical and
 R represents an optically pure enantiomer of
 a highly sterically hindered chiral hydrocarbon
 radical, preferably a branched alkyl or a cycloalkyl
 optionally substituted by one or more alkyl groups,
 10 is converted, so as to regio- and stereospecifically
 introduce the β -N-alkylamide and the α -hydroxyl or their
 cyclic precursors in a single stage by a Ritter
 reaction, which consists:

15 of the direct synthesis of a cyclic chain by
 reacting a cis- β -arylglycidate derivative of
 general formula I defined above with a nitrile of
 formula




20

in which


R', represents an aryl radical or a lower alkyl or
 lower perhaloalkyl radical, such as
 trichloromethyl,

in the presence of a Lewis acid or of a protonic acid, in anhydrous medium, in order to obtain the oxazoline of general formula IIb



in which Ar, R and R', are defined above.

5 2. Process according to one of claim 1, characterized in that the cis-β-arylglycidate derivative of general formula I



10 in which

Ar is defined as in claim 1 and
 R represents an optically pure enantiomer of a highly sterically hindered chiral hydrocarbon radical, preferably a branched alkyl or a cycloalkyl optionally substituted by one or more alkyl groups,
 is prepared by reacting the aldehyde of formula



15 with the haloacetate of formula



Ar and R being defined as in claim 1 and

X representing a halogen, in particular a chlorine or a bromine.

3. Process according to claim 1, characterized in that R represents an optically pure enantiomer of a highly sterically hindered chiral hydrocarbon radical, advantageously a cycloalkyl substituted by one or more alkyl groups, in particular a cyclohexyl.

4. Process according to claim 3, characterized in that R is one of the enantiomers of the menthyl radical, in particular (+)-menthyl.


5. Process according to one of claims 1 to 4, characterized in that the cis- β -phenylglycidate derivative of general formula I is of (2R,3R) configuration and the derivatives of general formula IIb obtained is of (2R,3S) configuration.

6. Process according to one of claims 1 to 5, characterized in that Ar and R₁ represent a phenyl.

7. Process according to claim 1 to 6, characterized in that the Lewis acid is chosen from the boron trifluoride acetic acid complex, boron trifluoride etherate, antimony pentachloride, tin tetrachloride or titanium tetrachloride and the protonic acid is tetrafluoroboric acid.

8. Process according to one of claims 1 to 7, characterized in that the derivatives of formula IIb defined as in claim 1 in which R represents a hydrogen atom are obtained by controlled saponification.

9. A compound of formula:



IIIb

in which:

Ar represents an aryl radical,

R represents an optically pure enantiomer of a highly sterically hindered chiral hydrocarbon radical, preferably a branched alkyl or a cycloalkyl optionally substituted by one or more alkyl groups and

R'₂ represents aryl radical above or a lower alkyl or lower perhaloalkyl radical, such as trichloromethyl.

לוצאטו את לוצאטו
LUZZATTO & LUZZATTO

By:

ע"י:



**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.